

Functional Equations of L -Functions for Symmetric Products of the Kloosterman Sheaf *

Lei Fu

Institute of Mathematics, Nankai University, Tianjin, P. R. China
leifu@nankai.edu.cn

Daqing Wan

Department of Mathematics, University of California, Irvine, CA 92697
dwan@math.uci.edu

Abstract

We determine the (arithmetic) local monodromy at 0 and at ∞ of the Kloosterman sheaf using local Fourier transformations and Laumon's stationary phase principle. We then calculate ϵ -factors for symmetric products of the Kloosterman sheaf. Using Laumon's product formula, we get functional equations of L -functions for these symmetric products, and prove a conjecture of Evans on signs of constants of functional equations.

Key words: Kloosterman sheaf, ϵ -factor, ℓ -adic Fourier transformation.

Mathematics Subject Classification: 11L05, 14G15.

Introduction

Let $p \neq 2$ be a prime number and let $\mathbb{F}_p$ be the finite field with p elements. Fix an algebraic closure $\mathbb{F}$ of $\mathbb{F}_p$. Denote the projective line over $\mathbb{F}_p$ by $\mathbb{P}^1$. For any power q of p , let $\mathbb{F}_q$ be the finite subfield of $\mathbb{F}$ with q elements. Let ℓ be a prime number different from p . Fix a nontrivial additive character $\psi : \mathbb{F}_p \rightarrow \overline{\mathbb{Q}}_\ell^*$. For any $x \in \mathbb{F}_q^*$, we define the one variable Kloosterman sum by

$$\text{Kl}_2(\mathbb{F}_q, x) = \sum_{\lambda \in \mathbb{F}_q^*} \psi \left(\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p} \left(\lambda + \frac{x}{\lambda} \right) \right).$$

In [3], Deligne constructs a lisse $\overline{\mathbb{Q}}_\ell$ -sheaf Kl_2 of rank 2 on $\mathbb{G}_m = \mathbb{P}^1 - \{0, \infty\}$, which we call the Kloosterman sheaf, such that for any $x \in \mathbb{G}_m(\mathbb{F}_q) = \mathbb{F}_q^*$, we have

$$\text{Tr}(F_x, \text{Kl}_{2, \bar{x}}) = -\text{Kl}_2(\mathbb{F}_q, x),$$

*The research of Lei Fu is supported by the NSFC (10525107).

where F_x is the geometric Frobenius element at the point x . For a positive integer k , the L -function $L(\mathbb{G}_m, \text{Sym}^k(\text{Kl}_2), T)$ of the k -th symmetric product of Kl_2 was first studied by Robba [15] via Dwork's p -adic methods. Motivated by applications in coding theory, by connections with modular forms, p -adic modular forms and Dwork's unit root zeta functions, there has been a great deal of recent interests to understand $L(\mathbb{G}_m, \text{Sym}^k(\text{Kl}_2), T)$ as much as possible for all k and for all p . This quickly raises a large number of interesting new problems.

Let $j : \mathbb{G}_m \rightarrow \mathbb{P}^1$ be the inclusion. We shall be interested in the L -function

$$M_k(p, T) := L(\mathbb{P}^1, j_*(\text{Sym}^k(\text{Kl}_2)), T).$$

This is the non-trivial factor of $L(\mathbb{G}_m, \text{Sym}^k(\text{Kl}_2), T)$. The trivial factor of $L(\mathbb{G}_m, \text{Sym}^k(\text{Kl}_2), T)$ was completely determined in Fu-Wan [6]. By general theory of Grothendieck-Deligne, the non-trivial factor $M_k(p, T)$ is a polynomial in T with integer coefficients, pure of weight $k + 1$. Its degree $\delta_k(p)$ can be easily extracted from Fu-Wan [7] Proposition 2.3, Lemmas 4.1 and 4.2:

$$\delta_k(p) = \begin{cases} \frac{k-1}{2} - \left\lfloor \frac{k}{2p} + \frac{1}{2} \right\rfloor & \text{if } k \text{ is odd,} \\ 2 \left(\left\lfloor \frac{k-2}{4} \right\rfloor - \left\lfloor \frac{k}{2p} \right\rfloor \right) & \text{if } k \text{ is even.} \end{cases}$$

For fixed k , the variation of $M_k(p, T)$ as p varies should be explained by an automorphic form, see Choi-Evans [2] and Evans [4] for the precise relations in the cases $k \leq 7$ and Fu-Wan [8] for a motivic interpretation for all k . For $k \leq 4$, the degree $\delta_k(p) \leq 1$ and $M_k(p, T)$ can be determined easily. For $k = 5$, the degree $\delta_5(p) = 2$ for $p > 5$. The quadratic polynomial $M_5(p, T)$ is explained by an explicit modular form [14]. For $k = 6$, the degree $\delta_6(p) = 2$ for $p > 6$. The quadratic polynomial $M_6(p, T)$ is again explained by an explicit modular form [9]. For $k = 7$, the degree $\delta_7(p) = 3$ for $p > 7$. The cubic polynomial $M_7(p, T)$ is conjecturally explained in a more subtle way by an explicit modular form in Evans [4]. We will return to this conjecture later in the introduction.

For fixed p , the variation of $M_k(p, T)$ as k varies p -adically should be related to p -adic automorphic forms and p -adic L -functions. No progress has been made along this direction. The p -adic limit of $M_k(p, T)$ as k varies p -adically links to an important example of Dwork's unit root zeta function, see the introduction in Wan [18]. The polynomial $M_k(p, T)$ can be used to determine the weight distribution of certain codes, see Moisisio [12][13], and this has been studied extensively for small p and small k . The p -adic Newton polygon (the p -adic slopes) of $M_k(p, T)$ remains largely mysterious.

By Katz [10] 4.1.11, we have $(\text{Kl}_2)^\vee = \text{Kl}_2 \otimes \overline{\mathbb{Q}_\ell}(1)$. So for any natural number k , we have

$$(\text{Sym}^k(\text{Kl}_2))^\vee = \text{Sym}^k(\text{Kl}_2) \otimes \overline{\mathbb{Q}_\ell}(k).$$

General theory (confer [11] 3.1.1) shows that $M_k(p, T)$ satisfies the functional equation

$$M_k(p, T) = cT^\delta M_k\left(p, \frac{1}{p^{k+1}T}\right),$$

where

$$\begin{aligned} c &= \prod_{i=0}^2 \det(-F, H^i(\mathbb{P}_{\mathbb{F}}^1, j_*(\text{Sym}^k(\text{Kl}_2)))^{(-1)^{i+1}}), \\ \delta &= -\chi(\mathbb{P}_{\mathbb{F}}^1, j_*(\text{Sym}^k(\text{Kl}_2))) = \delta_k(p), \end{aligned}$$

and F denotes the Frobenius correspondence. Applying the functional equation twice, we get

$$c^2 = p^{(k+1)\delta}.$$

Based on numerical computation, Evans [4] suggests that the sign of c should be $-\left(\frac{p}{105}\right)$ (the Jacobi symbol) for $k = 7$, and $-\left(\frac{p}{1155}\right)$ for $k = 11$. In this paper, we determine c for all k and all $p > 2$. The main result of this paper is the following theorem.

Theorem 0.1. *Let $p > 2$ be an odd prime. If k is even, we have*

$$c = p^{(k+1)(\lfloor \frac{k-2}{4} \rfloor - \lfloor \frac{k}{2p} \rfloor)}.$$

If k is odd, we have

$$c = (-1)^{\frac{k-1}{2} + \lfloor \frac{k}{2p} + \frac{1}{2} \rfloor} p^{\frac{k+1}{2}(\frac{k-1}{2} - \lfloor \frac{k}{2p} + \frac{1}{2} \rfloor)} \left(\frac{-2}{p}\right)^{\lfloor \frac{k}{2p} + \frac{1}{2} \rfloor} \prod_{j \in \{0, 1, \dots, \lfloor \frac{k}{2} \rfloor\}, p \nmid 2j+1} \left(\frac{(-1)^j(2j+1)}{p}\right).$$

Corollary 0.2. *If k is even and $p > 2$, the sign of c is always 1. If k is odd and $p > k$, the sign of c is*

$$(-1)^{\frac{k-1}{2}} \prod_{j \in \{0, 1, \dots, \lfloor \frac{k}{2} \rfloor\}, p \nmid 2j+1} \left(\frac{(-1)^j(2j+1)}{p}\right).$$

In the above corollary, if we take $k = 7$, we see that the sign of c for $p > 7$ is

$$-\left(\frac{1 \cdot (-3) \cdot 5 \cdot (-7)}{p}\right) = -\left(\frac{105}{p}\right) = -\left(\frac{p}{105}\right);$$

if we take $k = 11$, we see that the sign of c for $p > 11$ is

$$-\left(\frac{1 \cdot (-3) \cdot 5 \cdot (-7) \cdot 9 \cdot (-11)}{p}\right) = -\left(\frac{-1155}{p}\right) = -\left(\frac{p}{1155}\right),$$

consistent with Evans' calculation.

In the case $k = 7$, Evans proposed a precise description of $M_7(p, T)$ in terms of modular forms. For $k = 7$ and $p > 7$, the polynomial $M_7(p, T)$ has degree 3. Write

$$M_7(p, T) = 1 + a_p T + d_p T^2 + e_p T^3.$$

The functional equation and our sign determination show that one of the reciprocal roots for $M_7(p, T)$ is $(\frac{p}{105}) p^4$ and $e_p = -(\frac{p}{105}) p^{12}$. Denote the other two reciprocal roots by λ_p and μ_p which are Weil numbers of weight 8. We deduce that

$$a_p = -\left(\left(\frac{p}{105}\right) p^4 + \lambda_p + \mu_p\right), \quad \lambda_p \mu_p = p^8, \quad |\lambda_p| = |\mu_p| = p^4.$$

To explain the numerical calculation of Evans, Katz suggests that there exists a two dimensional representation

$$\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}(\overline{\mathbb{Q}}_\ell^2)$$

unramified for $p > 7$ and a Dirichlet character χ such that

$$\begin{aligned} \alpha_p^2 &= \chi(p) \left(\frac{p}{105}\right) \frac{\lambda_p}{p^4}, \\ \beta_p^2 &= \chi(p) \left(\frac{p}{105}\right) \frac{\mu_p}{p^4}, \\ \alpha_p \beta_p &= \chi(p), \end{aligned}$$

where α_p and β_p are the eigenvalues of the geometric Frobenius element at p under ρ . We then have

$$\begin{aligned} 1 - \left(\frac{p}{105}\right) \frac{a_p}{p^4} &= 2 + \left(\frac{p}{105}\right) \frac{\lambda_p}{p^4} + \left(\frac{p}{105}\right) \frac{\mu_p}{p^4} \\ &= \overline{\chi}(p)(2\alpha_p \beta_p + \alpha_p^2 + \beta_p^2) \\ &= \overline{\chi}(p)(\alpha_p + \beta_p)^2. \end{aligned}$$

Set $b(p) = p(\alpha_p + \beta_p)$. Evans [4] conjectured that $b(p)$ is the p -th Hecke eigenvalue for a weight 3 newform f on $\Gamma_0(525)$. Our a_p equals $-c_p p^2$ in [4].

Our proof of Theorem 0.1 naturally splits into two parts, corresponding to the two ramification points at 0 and ∞ . Let t be the coordinate of $\mathbb{A}^1 = \mathbb{P}^1 - \{\infty\}$. For any closed point x in $\mathbb{P}^1$, let $\mathbb{P}_{(x)}^1$ be the henselization of $\mathbb{P}^1$ at x . By Laumon's product formula [11] 3.2.1.1, we have

$$c = p^{k+1} \prod_{x \in |\mathbb{P}^1|} \epsilon(\mathbb{P}_{(x)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(x)}^1}, dt|_{\mathbb{P}_{(x)}^1}),$$

where $|\mathbb{P}^1|$ is the set of all closed points of $\mathbb{P}^1$. When $x \neq 0, \infty$, the sheaf $\text{Sym}^k(\text{Kl}_2)|_{\mathbb{P}^1_{(x)}}$ is lisse and the order of dt at x is 0. So by [11] 3.1.5.4 (ii) and (v), we have

$$\epsilon(\mathbb{P}^1_{(x)}, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}^1_{(x)}}, dt|_{\mathbb{P}^1_{(x)}}) = 1$$

for $x \neq 0, \infty$. Therefore

$$c = p^{k+1} \epsilon(\mathbb{P}^1_{(0)}, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}^1_{(0)}}, dt|_{\mathbb{P}^1_{(0)}}) \epsilon(\mathbb{P}^1_{(\infty)}, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}^1_{(\infty)}}, dt|_{\mathbb{P}^1_{(\infty)}}).$$

In §1, we prove the following.

Proposition 0.3. *We have*

$$\epsilon(\mathbb{P}^1_{(0)}, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}^1_{(0)}}, dt|_{\mathbb{P}^1_{(0)}}) = (-1)^k p^{\frac{k(k+1)}{2}}.$$

In §2, we prove the following.

Proposition 0.4. $\epsilon(\mathbb{P}^1_{(\infty)}, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}^1_{(\infty)}}, dt|_{\mathbb{P}^1_{(\infty)}})$ equals

$$p^{-(k+1)(\frac{k+8}{4} + [\frac{k}{2p}])}$$

if $k = 2r$ for an even r ,

$$p^{-(k+1)(\frac{k+6}{4} + [\frac{k}{2p}])}$$

if $k = 2r$ for an odd r , and

$$(-1)^{\frac{k+1}{2} + [\frac{k}{p}] - [\frac{k}{2p}]} p^{-\frac{k+1}{2}(\frac{k+5}{2} + [\frac{k}{p}] - [\frac{k}{2p}])} \left(\frac{-2}{p}\right)^{[\frac{k}{p}] - [\frac{k}{2p}]} \prod_{j \in \{0, 1, \dots, [\frac{k}{2}]\}, p \nmid 2j+1} \left(\frac{(-1)^j (2j+1)}{p}\right)$$

if $k = 2r + 1$.

We deduce from the above two propositions the constant c as stated in Theorem 0.1 using the following facts:

$$\left\lceil \frac{k-2}{4} \right\rceil = \begin{cases} \frac{k-4}{4} & \text{if } k = 2r \text{ for an even } r, \\ \frac{k-2}{4} & \text{if } k = 2r \text{ for an odd } r, \end{cases}$$

$$\left\lceil \frac{k}{p} \right\rceil - \left\lceil \frac{k}{2p} \right\rceil = \left\lceil \frac{k}{2p} + \frac{1}{2} \right\rceil \text{ if } k \text{ is odd.}$$

To get Proposition 0.4, we first have to determine the local (arithmetic) monodromy of Kl_2 at ∞ . This is Theorem 2.1 in §2, which is of interest itself, and is proved by using local Fourier transformations and Laumon's stationary phase principle.

1 Calculation of $\epsilon(\mathbb{P}_{(0)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(0)}^1}, dt|_{\mathbb{P}_{(0)}^1})$

Let η_0 be the generic point of $\mathbb{P}_{(0)}^1$, let $\bar{\eta}_0$ be a geometric point located at η_0 , and let V be an $\overline{\mathbb{Q}}_\ell$ -representation of $\text{Gal}(\bar{\eta}_0/\eta_0)$. Suppose the inertia subgroup I_0 of $\text{Gal}(\bar{\eta}_0/\eta_0)$ acts unipotently on V . Consider the ℓ -adic part of the cyclotomic character

$$t_\ell : I_0 \rightarrow \mathbb{Z}_\ell(1), \sigma \mapsto \left(\frac{\sigma(\sqrt[n]{t})}{\sqrt[n]{t}} \right).$$

Note that for any σ in the inertia subgroup, the ℓ^n -th root of unity $\frac{\sigma(\sqrt[n]{t})}{\sqrt[n]{t}}$ does not depend on the choice of the ℓ^n -th root $\sqrt[n]{t}$ of t . Since I_0 acts on V unipotently, there exists a nilpotent homomorphism

$$N : V(1) \rightarrow V$$

such that the action of $\sigma \in I_0$ on V is given by $\exp(t_\ell(\sigma).N)$. Fix a lifting $F \in \text{Gal}(\bar{\eta}_0/\eta_0)$ of the geometric Frobenius element in $\text{Gal}(\mathbb{F}/\mathbb{F}_p)$.

Lemma 1.1. *Notation as above. Let $V = \text{Kl}_{2, \bar{\eta}_0}$. There exists a basis $\{e_0, e_1\}$ of V such that*

$$F(e_0) = e_0, F(e_1) = pe_1$$

$$N(e_0) = 0, N(e_1) = e_0.$$

Proof. This is the $n = 2$ case of Proposition 1.1 in [6]. □

Lemma 1.2. *Keep the notation in Lemma 1.1. Let $\{f_0, \dots, f_k\}$ be the basis of $\text{Sym}^k(V) = \text{Sym}^k(\text{Kl}_{2, \bar{\eta}_0})$ defined by $f_i = \frac{1}{i!} e_0^{k-i} e_1^i$. We have*

$$F(f_i) = p^i f_i, N(f_i) = f_{i-1},$$

where we regard f_{i-1} as 0 if $i = 0$.

Proof. Use the fact that for any $v_1, \dots, v_k \in V$, we have the following identities in $\text{Sym}^k(V)$:

$$\begin{aligned} F(v_1 \cdots v_k) &= F(v_1) \cdots F(v_k), \\ N(v_1 \cdots v_k) &= \sum_{i=1}^k v_1 \cdots v_{i-1} N(v_i) v_{i+1} \cdots v_k. \end{aligned}$$

□

Corollary 1.3. *The sheaf $\mathrm{Sym}^k(\mathrm{Kl}_2)|_{\eta_0}$ has a filtration*

$$0 = \mathcal{F}_{-1} \subset \mathcal{F}_0 \subset \cdots \subset \mathcal{F}_k = \mathrm{Sym}^k(\mathrm{Kl}_2)|_{\eta_0}$$

such that

$$\mathcal{F}_i/\mathcal{F}_{i-1} \cong \overline{\mathbb{Q}}_l(-i)$$

for any $i = 0, \dots, k$.

Proof. This follows from Lemma 1.2 by taking $\mathcal{F}_i$ to be the sheaf on η_0 corresponding to the galois representation $\mathrm{Span}(f_0, \dots, f_i)$ of $\mathrm{Gal}(\bar{\eta}_0/\eta_0)$. $\square$

The following is Proposition 0.3 in the introduction.

Proposition 1.4. *We have*

$$\epsilon(\mathbb{P}_{(0)}^1, j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}, dt|_{\mathbb{P}_{(0)}^1}) = (-1)^k p^{\frac{k(k+1)}{2}}.$$

Proof. Let $u : \eta_0 \rightarrow \mathbb{P}_{(0)}^1$ and $v : \{0\} \rightarrow \mathbb{P}_{(0)}^1$ be the immersions. By [11] 3.1.5.4 (iii) and (v), we have

$$\begin{aligned} \epsilon(\mathbb{P}_{(0)}^1, u_* \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1}) &= 1, \\ \epsilon(\mathbb{P}_{(0)}^1, v_* \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1}) &= \det(-F_0, \overline{\mathbb{Q}}_\ell(-i))^{-1} = -\frac{1}{p^i}. \end{aligned}$$

We have an exact sequence

$$0 \rightarrow u_! \overline{\mathbb{Q}}_\ell(-i) \rightarrow u_* \overline{\mathbb{Q}}_\ell(-i) \rightarrow v_* \overline{\mathbb{Q}}_\ell(-i) \rightarrow 0.$$

It follows from [11] 3.1.5.4 (ii) that we have

$$\begin{aligned} \epsilon(\mathbb{P}_{(0)}^1, u_! \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1}) &= \frac{\epsilon(\mathbb{P}_{(0)}^1, u_* \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1})}{\epsilon(\mathbb{P}_{(0)}^1, v_* \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1})} \\ &= -p^i. \end{aligned}$$

By Corollary 1.3 and [11] 3.1.5.4 (ii), we have

$$\begin{aligned} \epsilon(\mathbb{P}_{(0)}^1, j_!(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}, dt|_{\mathbb{P}_{(0)}^1}) &= \prod_{i=0}^k \epsilon(\mathbb{P}_{(0)}^1, u_!(\mathcal{F}_i/\mathcal{F}_{i+1}), dt|_{\mathbb{P}_{(0)}^1}) \\ &= \prod_{i=0}^k \epsilon(\mathbb{P}_{(0)}^1, u_! \overline{\mathbb{Q}}_\ell(-i), dt|_{\mathbb{P}_{(0)}^1}) \\ &= \prod_{i=0}^k (-p^i). \end{aligned}$$

Moreover, by Lemma 1.2, we have

$$v^*(j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}) \cong \overline{\mathbb{Q}}_\ell,$$

and hence

$$\epsilon(\mathbb{P}_{(0)}^1, v_* v^*(j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}), dt|_{\mathbb{P}_{(0)}^1}) = -1.$$

So we have

$$\begin{aligned} & \epsilon(\mathbb{P}_{(0)}^1, j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}, dt|_{\mathbb{P}_{(0)}^1}) \\ &= \epsilon(\mathbb{P}_{(0)}^1, j_!(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}, dt|_{\mathbb{P}_{(0)}^1}) \epsilon(\mathbb{P}_{(0)}^1, v_* v^*(j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(0)}^1}), dt|_{\mathbb{P}_{(0)}^1}) \\ &= \prod_{i=1}^k (-p^i) \\ &= (-1)^k p^{\frac{k(k+1)}{2}}. \end{aligned}$$

□

2 Calculation of $\epsilon(\mathbb{P}_{(\infty)}^1, j_*(\mathrm{Sym}^k(\mathrm{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1})$

We first introduce some notations. Fix a nontrivial additive character $\psi : \mathbb{F}_p \rightarrow \overline{\mathbb{Q}}_\ell^*$ and define Kl_2 as in the introduction. Fix a separable closure $\overline{\mathbb{F}_p(t)}$ of $\mathbb{F}_p(t)$. Let x be an element in $\overline{\mathbb{F}_p(t)}$ satisfying $x^p - x = t$. Then $\mathbb{F}_p(t, x)$ is galois over $\mathbb{F}_p(t)$. We have a canonical isomorphism

$$\mathbb{F}_p \xrightarrow{\cong} \mathrm{Gal}(\mathbb{F}_p(t, x)/\mathbb{F}_p(t))$$

which sends each $a \in \mathbb{F}_p$ to the element in $\mathrm{Gal}(\mathbb{F}_p(t, x)/\mathbb{F}_p(t))$ defined by $x \mapsto x + a$. Let $\mathcal{L}_\psi$ be the galois representation defined by

$$\mathrm{Gal}(\overline{\mathbb{F}_p(t)}/\mathbb{F}_p(t)) \rightarrow \mathrm{Gal}(\mathbb{F}_p(t, x)/\mathbb{F}_p(t)) \xrightarrow{\cong} \mathbb{F}_p \xrightarrow{\psi^{-1}} \overline{\mathbb{Q}}_\ell^*.$$

It is unramified outside ∞ and totally wild at ∞ with Swan conductor 1. This galois representation defines a lisse $\overline{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{A}^1$ which we still denote by $\mathcal{L}_\psi$. Let X be an $\mathbb{F}_p$ -scheme. Any section f in $\mathcal{O}_X(X)$ defines an $\mathbb{F}_p$ -algebra homomorphism

$$\mathbb{F}_p[t] \rightarrow \mathcal{O}_X(X), \quad t \mapsto f,$$

and hence an $\mathbb{F}_p$ -morphism of schemes

$$f : X \rightarrow \mathbb{A}^1.$$

We denote the lisse $\overline{\mathbb{Q}_\ell}$ -sheaf $f^*\mathcal{L}_\psi$ on X by $\mathcal{L}_\psi(f)$. For any $f_1, f_2 \in \mathcal{O}_X(X)$, we have

$$\mathcal{L}_\psi(f_1) \otimes \mathcal{L}_\psi(f_2) \cong \mathcal{L}_\psi(f_1 + f_2).$$

Recall that $p \neq 2$. Let y be an element in $\overline{\mathbb{F}_p(t)}$ satisfying $y^2 = t$. Then $\mathbb{F}_p(t, y)$ is galois over $\mathbb{F}_p(t)$. We have a canonical isomorphism

$$\{\pm 1\} \xrightarrow{\cong} \text{Gal}(\mathbb{F}_p(t, y)/\mathbb{F}_p(t))$$

which sends -1 to the element in $\text{Gal}(\mathbb{F}_p(t, y)/\mathbb{F}_p(t))$ defined by $y \mapsto -y$. Let

$$\chi : \{\pm 1\} \rightarrow \overline{\mathbb{Q}_\ell}^*$$

be the (unique) nontrivial character. Define $\mathcal{L}_\chi$ to be the galois representation defined by

$$\text{Gal}(\overline{\mathbb{F}_p(t)}/\mathbb{F}_p(t)) \rightarrow \text{Gal}(\mathbb{F}_p(t, y)/\mathbb{F}_p(t)) \xrightarrow{\cong} \{\pm 1\} \xrightarrow{\chi^{-1}} \overline{\mathbb{Q}_\ell}^*.$$

It is unramified outside 0 and ∞ , and tamely ramified at 0 and ∞ . This galois representation defines a lisse $\overline{\mathbb{Q}_\ell}$ -sheaf on $\mathbb{G}_m$ which we still denote by $\mathcal{L}_\chi$.

Let $\theta : \text{Gal}(\mathbb{F}/\mathbb{F}_p) \rightarrow \overline{\mathbb{Q}_\ell}^*$ be a character of the galois group of the finite field. Denote by $\mathcal{L}_\theta$ the galois representation

$$\text{Gal}(\overline{\mathbb{F}_p(t)}/\mathbb{F}_p(t)) \rightarrow \text{Gal}(\mathbb{F}/\mathbb{F}_p) \xrightarrow{\theta} \overline{\mathbb{Q}_\ell}^*.$$

It is unramified everywhere, and hence defines a lisse $\overline{\mathbb{Q}_\ell}$ -sheaf on $\mathbb{P}^1$ which we still denote by $\mathcal{L}_\theta$.

Theorem 2.1. *Notation as above. Let η_∞ be the generic point of $\mathbb{P}_{(\infty)}^1$. Then $\text{Kl}_2|_{\eta_\infty}$ is isomorphic to the restriction to η_∞ of the sheaf*

$$[2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_0},$$

where $[2] : \mathbb{G}_m \rightarrow \mathbb{G}_m$ is the morphism defined by $x \mapsto x^2$, and

$$\theta_0 : \text{Gal}(\mathbb{F}/\mathbb{F}_p) \rightarrow \overline{\mathbb{Q}_\ell}^*$$

is the character sending the geometric Frobenius element F in $\text{Gal}(\mathbb{F}/\mathbb{F}_p)$ to the Gauss sum

$$\theta_0(F) = g(\chi, \psi) = - \sum_{x \in \mathbb{F}_p^*} \left(\frac{x}{p} \right) \psi(x).$$

Proof. By [8] Proposition 1.1, we have

$$\mathrm{Kl}_2 = \mathcal{F} \left(j! \mathcal{L}_\psi \left(\frac{1}{t} \right) \right) |_{\mathbb{G}_m}, \quad (1)$$

where $\mathcal{F}$ is the ℓ -adic Fourier transformation and $j : \mathbb{G}_m \rightarrow \mathbb{A}^1$ is the inclusion. Let

$$\pi_1, \pi_2 : \mathbb{G}_m \times_{\mathbb{F}_p} \mathbb{G}_m \rightarrow \mathbb{G}_m$$

be the projections. Using the proper base change theorem and the projection formula ([1] XVII 5.2.6 and 5.2.9), one can verify

$$[2]^* \left(\mathcal{F} \left(j! \mathcal{L}_\psi \left(\frac{1}{t} \right) \right) |_{\mathbb{G}_m} \right) \cong R\pi_{2!} \left(\mathcal{L}_\psi \left(\frac{1}{t} + tt'^2 \right) \right) [1], \quad (2)$$

where

$$\frac{1}{t} + tt'^2 : \mathbb{G}_m \times_{\mathbb{F}_p} \mathbb{G}_m \rightarrow \mathbb{A}^1$$

is the morphism corresponding to the $\mathbb{F}_p$ -algebra homomorphism

$$\mathbb{F}_p[t] \rightarrow \mathbb{F}_p[t, 1/t, t', 1/t'], \quad t \mapsto \frac{1}{t} + tt'^2.$$

Consider the isomorphism

$$\tau : \mathbb{G}_m \times_{\mathbb{F}_p} \mathbb{G}_m \rightarrow \mathbb{G}_m \times_{\mathbb{F}_p} \mathbb{G}_m, \quad (t, t') \mapsto \left(\frac{t}{t'}, t' \right).$$

We have $\pi_2 \tau = \pi_2$. So

$$R\pi_{2!} \left(\mathcal{L}_\psi \left(\frac{1}{t} + tt'^2 \right) \right) \cong R(\pi_2 \tau)! \tau^* \left(\mathcal{L}_\psi \left(\frac{1}{t} + tt'^2 \right) \right) \cong R\pi_{2!} \mathcal{L}_\psi \left(\left(\frac{1}{t} + t \right) t' \right). \quad (3)$$

Consider the morphism

$$g : \mathbb{G}_m \rightarrow \mathbb{A}^1, \quad t \mapsto \frac{1}{t} + t.$$

Again using the proper base change theorem and the projection formula, one can verify

$$\mathcal{F}(Rg! \overline{\mathbb{Q}}_\ell) \cong R\pi_{2!} \mathcal{L}_\psi \left(\left(\frac{1}{t} + t \right) t' \right) [1]. \quad (4)$$

From the isomorphisms (1)-(4), we get

$$[2]^* \mathrm{Kl}_2 \cong \mathcal{F}(Rg! \overline{\mathbb{Q}}_\ell) |_{\mathbb{G}_m}.$$

By Lemma 2.2 below, the stationary phase principle of Laumon [11] 2.3.3.1 (iii), and [11] 2.5.3.1, as representations of $\text{Gal}(\bar{\eta}_{\infty'}/\eta_{\infty'})$, we have

$$\begin{aligned}\mathcal{H}^0(\mathcal{F}(Rg_!\bar{\mathbb{Q}}_\ell))_{\bar{\eta}_{\infty'}} &\cong \mathcal{F}^{(2,\infty')}(\mathcal{L}_\chi) \bigoplus \mathcal{F}^{(-2,\infty')}(\mathcal{L}_\chi) \\ &\cong (\mathcal{L}_\psi(2t') \otimes \mathcal{F}^{(0,\infty')}(\mathcal{L}_\chi)) \bigoplus (\mathcal{L}_\psi(-2t') \otimes \mathcal{F}^{(0,\infty')}(\mathcal{L}_\chi)) \\ &\cong (\mathcal{L}_\psi(2t') \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0}) \bigoplus (\mathcal{L}_\psi(-2t') \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0}).\end{aligned}$$

Hence

$$([2]^*\text{Kl}_2)|_{\eta_\infty} \cong (\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0})|_{\eta_\infty} \bigoplus (\mathcal{L}_\psi(-2t) \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0})|_{\eta_\infty}.$$

Note that this decomposition of $([2]^*\text{Kl}_2)|_{\eta_\infty}$ is non-isotypical. By [16] Proposition 24 on p. 61, and the fact that $\text{Kl}_2|_{\eta_\infty}$ is irreducible (since its Swan conductor is 1), we have

$$\text{Kl}_2|_{\eta_\infty} \cong [2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0})|_{\eta_\infty}.$$

We have

$$\begin{aligned}[2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi \otimes \mathcal{L}_{\theta_0}) &\cong [2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi \otimes [2]^*\mathcal{L}_{\theta_0}) \\ &\cong [2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_0}.\end{aligned}$$

Here we use the fact that $[2]^*\mathcal{L}_{\theta_0} \cong \mathcal{L}_{\theta_0}$. Hence

$$\text{Kl}_2|_{\eta_\infty} \cong \left([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_0}\right)|_{\eta_\infty}.$$

□

Lemma 2.2. *For the morphism*

$$g : \mathbb{G}_m \rightarrow \mathbb{A}^1, \quad t \mapsto \frac{1}{t} + t,$$

the following holds:

- (i) $Rg_!\bar{\mathbb{Q}}_\ell$ is a $\bar{\mathbb{Q}}_\ell$ -sheaf on $\mathbb{A}^1$ which is lisse outside the rational points 2 and -2 .
- (ii) $Rg_!\bar{\mathbb{Q}}_\ell$ is unramified at ∞ .
- (iii) Let P be one of the rational points 2 or -2 , and let $\tilde{\mathbb{A}}_{(P)}^1$ be the henselization of $\mathbb{A}^1$ at P .

We have

$$(Rg_!\bar{\mathbb{Q}}_\ell)|_{\tilde{\mathbb{A}}_{(P)}^1} \cong \bar{\mathbb{Q}}_\ell \oplus \mathcal{L}_{\chi,!},$$

where $\mathcal{L}_{\chi,!}$ denotes the extension by 0 of the Kummer sheaf $\mathcal{L}_\chi$ on the generic point of $\tilde{\mathbb{A}}_{(P)}^1$ to $\tilde{\mathbb{A}}_{(P)}^1$.

Proof. We have

$$\frac{\partial g}{\partial t} = -\frac{1}{t^2} + 1.$$

So $\frac{\partial g}{\partial t}$ vanishes at the points $t = \pm 1$. We have

$$\begin{aligned} g(\pm 1) &= \pm 2, \\ \frac{\partial^2 g}{\partial t^2}(\pm 1) &= \pm 2 \neq 0. \end{aligned}$$

It follows that g is tamely ramified above ± 2 with ramification index 2, and g is étale elsewhere.

Consider the morphism

$$\bar{g} : \mathbb{P}^1 \rightarrow \mathbb{P}^1, [t_0 : t_1] \mapsto [t_0 t_1 : t_0^2 + t_1^2].$$

We have $\bar{g}^{-1}(\infty) = \{0, \infty\}$. Hence

$$\bar{g}^{-1}(\mathbb{A}^1) = \mathbb{G}_m.$$

It is clear that

$$\bar{g}|_{\mathbb{G}_m} = g.$$

So $g : \mathbb{G}_m \rightarrow \mathbb{A}^1$ is a finite morphism of degree 2. Near 0, the morphism $\bar{g}$ can be expressed as

$$t \mapsto \frac{t}{1+t^2}.$$

Hence $\bar{g}$ is unramified at 0. Similarly $\bar{g}$ is also unramified at ∞ . Our lemma follows from these facts. $\square$

Remark 2.3. The first attempt to determine the monodromy at ∞ of the $(n-1)$ -variable Kloosterman sheaf $\text{Kl}_n|_{\eta_\infty}$ is done in Fu-Wan [7] Theorem 1.1, where we deduce from Katz [10] that

$$\text{Kl}_n|_{\eta_\infty} \cong \left([n]_* (\mathcal{L}_\psi(nt) \otimes \mathcal{L}_{\chi^{n-1}}) \otimes \mathcal{L}_\theta \otimes \overline{\mathbb{Q}}_\ell \left(\frac{1-n}{2} \right) \right) |_{\eta_\infty}$$

for some character $\theta : \text{Gal}(\mathbb{F}/\mathbb{F}_p) \rightarrow \overline{\mathbb{Q}}_\ell^*$, and an explicit description of θ^2 is given. Using induction on n , [8] Proposition 1.1, and adapting the argument in [5] to non-algebraically closed ground field, we can get an explicit description of θ . See [5] where the monodromy of the more general hypergeometric sheaf is treated (over algebraically closed field).

Lemma 2.4. *Keep the notation in Theorem 2.1. Let*

$$\theta_1 : \text{Gal}(\mathbb{F}/\mathbb{F}_p) \rightarrow \overline{\mathbb{Q}}_\ell^*$$

be the character defined by

$$\theta_1(\sigma) = \chi \left(\frac{\sigma(\sqrt{-1})}{\sqrt{-1}} \right)$$

for any $\sigma \in \text{Gal}(\mathbb{F}/\mathbb{F}_p)$. Note that the above expression is independent of the choice of the square root $\sqrt{-1} \in \mathbb{F}$ of -1 .

(i) If $k = 2r$ is even, $\text{Sym}^k(\text{Kl}_2)|_{\eta_\infty}$ is isomorphic to the restriction to η_∞ of the sheaf

$$\left(\mathcal{L}_{\chi^r} \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^r} \right) \oplus \left(\bigoplus_{i=0}^{r-1} [2]_* \mathcal{L}_\psi((4i-4r)t) \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^i} \right).$$

(ii) If $k = 2r + 1$ is odd, $\text{Sym}^k(\text{Kl}_2)|_{\eta_\infty}$ is isomorphic to the restriction to η_∞ of the sheaf

$$\bigoplus_{i=0}^r [2]_* (\mathcal{L}_\psi((4i-4r-2)t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_0^{2r+1}\theta_1^{i+1}}.$$

Proof. By Theorem 2.1, it suffices to calculate the restriction to η_∞ of $\text{Sym}^k([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi))$.

Let y, z, w be elements in $\overline{\mathbb{F}_p(t)}$ satisfying

$$y^2 = t, \quad z^p - z = y, \quad w^2 = y.$$

Fix a square root $\sqrt{-1}$ of -1 in $\mathbb{F}$. Then $\mathbb{F}_p(z, w, \sqrt{-1})$ and $\mathbb{F}_p(y)$ are galois extensions of $\mathbb{F}_p(t)$.

Let $G = \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(t))$ and $H = \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(y))$. Then H is normal in G , and we have canonical isomorphisms

$$G/H \xrightarrow{\cong} \text{Gal}(\mathbb{F}_p(y)/\mathbb{F}_p(t)) \xrightarrow{\cong} \{\pm 1\}.$$

Consider the case where $\sqrt{-1}$ does not lie in $\mathbb{F}_p$. We have an isomorphism

$$\mathbb{F}_p \times \{\pm 1\} \times \{\pm 1\} \xrightarrow{\cong} H = \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(y))$$

which maps $(a, \mu', \mu'') \in \mathbb{F}_p \times \{\pm 1\} \times \{\pm 1\}$ to the element $g_{(a, \mu', \mu'')} \in \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(y))$ defined by

$$g_{(a, \mu', \mu'')}(z) = z + a, \quad g_{(a, \mu', \mu'')}(w) = \mu' w, \quad g_{(a, \mu', \mu'')}(w) = \mu'' \sqrt{-1}.$$

(In the case where $\sqrt{-1}$ lies in $\mathbb{F}_p$, we have $\mathbb{F}_p(z, w, \sqrt{-1}) = \mathbb{F}_p(z, w)$, and we have an isomorphism

$$\mathbb{F}_p \times \{\pm 1\} \xrightarrow{\cong} H = \text{Gal}(\mathbb{F}_p(z, w)/\mathbb{F}_p(y))$$

which maps $(a, \mu) \in \mathbb{F}_p \times \{\pm 1\}$ to the element $g_{(a, \mu)} \in \text{Gal}(\mathbb{F}_p(z, w)/\mathbb{F}_p(y))$ defined by

$$g_{(a, \mu)}(z) = z + a, \quad g_{(a, \mu)}(w) = \mu w.$$

The following argument works for this case with slight modification. We leave to the reader to treat this case.) Let V be a one dimensional $\overline{\mathbb{Q}}_\ell$ -vector space with a basis e_0 . Define an action of H on V by

$$g_{(a,\mu',\mu'')}(e_0) = \psi(-2a)\chi(\mu'^{-1})e_0.$$

Then $[2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi)$ is just the composition of $\text{Ind}_H^G(V)$ with the canonical homomorphism

$$\text{Gal}(\overline{\mathbb{F}_p(t)}/\mathbb{F}_p(t)) \rightarrow \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(t)) = G.$$

Let g be the element in $G = \text{Gal}(\mathbb{F}_p(z, w, \sqrt{-1})/\mathbb{F}_p(t))$ defined by

$$g(z) = -z, \quad g(w) = \sqrt{-1}w, \quad g(\sqrt{-1}) = \sqrt{-1}.$$

Then the image of g in G/H is a generator of the cyclic group G/H . So G is generated by $g_{(a,\mu',\mu'')} \in H$ ($(a, \mu', \mu'') \in \mathbb{F}_p \times \{\pm 1\} \times \{\pm 1\}$) and g . The space $\text{Ind}_H^G(V)$ has a basis $\{e_0, e_1\}$ with

$$\begin{aligned} g(e_0) &= e_1, \\ g_{(a,\mu',\mu'')}(e_0) &= \psi(-2a)\chi(\mu'^{-1})e_0, \\ g_{(a,\mu',\mu'')}(e_1) &= \psi(2a)\chi(\mu'^{-1}\mu''^{-1})e_1, \\ g(e_1) &= g^2(e_0) = g_{(0,-1,1)}(e_0) = -e_0. \end{aligned}$$

Suppose $k = 2r$ is even. $\text{Sym}^k(\text{Ind}_H^G(V))$ has a basis

$$\{e_1^k, g(e_1^k), e_0e_1^{k-1}, g(e_0e_1^{k-1}), \dots, e_0^{r-1}e_1^{r+1}, g(e_0^{r-1}e_1^{r+1}), e_0^r e_1^r\},$$

and for each $i = 0, 1, \dots, r$, we have

$$\begin{aligned} g_{(a,\mu',\mu'')}(e_0^i e_1^{k-i}) &= \psi(-2ia)\chi(\mu'^{-i})\psi(2(k-i)a)\chi(\mu'^{-(k-i)}\mu''^{-(k-i)})e_0^i e_1^{k-i} \\ &= \psi(2(k-2i)a)\chi(\mu'^{-k})\chi(\mu''^{-(k-i)})e_0^i e_1^{k-i}. \end{aligned}$$

Using the fact that k is even and $\chi^2 = 1$, we get

$$g_{(a,\mu',\mu'')}(e_0^i e_1^{k-i}) = \psi(2(k-2i)a)\chi(\mu''^i)e_0^i e_1^{k-i}.$$

In particular, we have

$$g_{(a,\mu',\mu'')}(e_0^r e_1^r) = \chi(\mu''^r)e_0^r e_1^r.$$

Moreover, we have

$$g(e_0^r e_1^r) = e_1^r(g(e_1^r)) = (-1)^r e_0^r e_1^r.$$

It follows that

$$\mathrm{Sym}^k([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi)) \cong (\mathcal{L}_{\chi^r} \otimes \mathcal{L}_{\theta_1^r}) \oplus \left(\bigoplus_{i=0}^{r-1} [2]_*(\mathcal{L}_\psi(2(2i-k)t) \otimes \mathcal{L}_{\theta_1^i}) \right).$$

We have

$$[2]_*(\mathcal{L}_\psi(2(2i-k)t) \otimes \mathcal{L}_{\theta_1^i}) \cong [2]_*(\mathcal{L}_\psi(2(2i-k)t) \otimes [2]^* \mathcal{L}_{\theta_1^i}) \cong [2]_* \mathcal{L}_\psi(2(2i-k)t) \otimes \mathcal{L}_{\theta_1^i}.$$

So we have

$$\mathrm{Sym}^k([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi)) \cong (\mathcal{L}_{\chi^r} \otimes \mathcal{L}_{\theta_1^r}) \oplus \left(\bigoplus_{i=0}^{r-1} [2]_* \mathcal{L}_\psi(2(2i-k)t) \otimes \mathcal{L}_{\theta_1^i} \right).$$

Suppose $n = 2r + 1$ is odd. $\mathrm{Sym}^k(\mathrm{Ind}_H^G(V))$ has a basis

$$\{e_1^k, g(e_1^k), e_0 e_1^{k-1}, g(e_0 e_1^{k-1}), \dots, e_0^r e_1^{r+1}, g(e_0^r e_1^{r+1})\}.$$

Using the same calculation as above, we get

$$\mathrm{Sym}^k([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi)) \cong \bigoplus_{i=0}^r [2]_*(\mathcal{L}_\psi(2(2i-k)t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_1^{i+1}}.$$

Lemma 2.4 follows by twisting the above expressions of $\mathrm{Sym}^k([2]_*(\mathcal{L}_\psi(2t) \otimes \mathcal{L}_\chi))$ by $\mathcal{L}_{\theta_0^k}$. $\square$

Lemma 2.5. *Assume $a \in \mathbb{F}_p$ is nonzero. We have the following identities.*

- (i) $\epsilon(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_\ell, dt|_{\mathbb{P}_{(\infty)}^1}) = \frac{1}{p^2}.$
- (ii) $\epsilon(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_\ell, dt^2|_{\mathbb{P}_{(\infty)}^1}) = \frac{1}{p^3}.$
- (iii) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = -\frac{g(\chi, \psi)}{p^2}.$
- (iv) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}) = -\frac{g(\chi, \psi)}{p^3} \left(\frac{-2}{p} \right).$
- (v) $\epsilon(\mathbb{P}_{(\infty)}^1, j_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}) = \frac{1}{p^2} \left(\frac{2a}{p} \right).$
- (vi) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\psi(at)|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}) = \frac{1}{p^2}.$
- (vii) $\epsilon(\mathbb{P}_{(\infty)}^1, [2]_* \overline{\mathbb{Q}}_\ell|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = -\frac{g(\chi, \psi)}{p^4}.$
- (viii) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* [2]_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = -\frac{g(\chi, \psi)}{p^3} \left(\frac{2a}{p} \right).$
- (ix) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* [2]_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = \frac{g(\chi, \psi)^2}{p^4} \left(\frac{-2}{p} \right).$
- (x) $\epsilon(\mathbb{P}_{(\infty)}^1, j_* [2]_* \mathcal{L}_\psi(at)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = -\frac{g(\chi, \psi)}{p^3}.$

Proof. Let K_∞ be the completion of the field $k(\eta_\infty)$, let $\mathcal{O}_\infty$ be the ring of integers in K_∞ , and let $s = \frac{1}{t}$. Then s is a uniformizer of K_∞ . Denote the inclusion $\eta_\infty \rightarrow \mathbb{P}_{(\infty)}^1$ also by j . Let V be a $\overline{\mathbb{Q}_\ell}$ -sheaf of rank 1 on η_∞ , and let $\phi : K_\infty^* \rightarrow \overline{\mathbb{Q}_\ell}^*$ be the character corresponding to V via the reciprocity law. The Artin conductor $a(\phi)$ of ϕ is defined to be the smallest integer m such that $\phi|_{1+s^m\mathcal{O}_\infty} = 1$. For any nonzero meromorphic differential 1-form $\omega = f ds$ on $\mathbb{P}_{(\infty)}^1$, define the order $v_\infty(\omega)$ of ω to be the valuation $v_\infty(f)$ of f . By [11] 3.1.5.4 (v), we have

$$\epsilon(\mathbb{P}_{(\infty)}^1, j_* V, \omega) = \begin{cases} \phi(s^{v_\infty(\omega)}) p^{v_\infty(\omega)} & \text{if } \phi|_{\mathcal{O}_\infty^*} = 1, \\ \int_{s^{-(a(\phi)+v_\infty(\omega))}\mathcal{O}_\infty^*} \phi^{-1}(z) \psi(\text{Res}_\infty(z\omega)) dz & \text{if } \phi|_{\mathcal{O}_\infty^*} \neq 1, \end{cases}$$

where Res_∞ denotes the residue of a meromorphic 1-form at ∞ , and the integral is taken with respect to the Haar measure dz on K_∞ normalized by $\int_{\mathcal{O}_\infty} dz = 1$.

Note that $dt = -\frac{ds}{s^2}$ has order -2 at ∞ and $dt^2 = -\frac{2ds}{s^3}$ has order -3 . Applying the first case of the above formula for the ϵ -factor, we get (i) and (ii).

(iii) Taking $a = t = \frac{1}{s}$ and $b = z$ in the explicit reciprocity law in [17] XIV §3 Proposition 8, we see the character

$$\chi' : K_\infty^* \rightarrow \overline{\mathbb{Q}_\ell}^*$$

corresponding to $\mathcal{L}_\chi$ is given by

$$\chi'(z) = \chi^{-1} \left(\left(\frac{\bar{c}}{p} \right) \right),$$

where

$$c = (-1)^{-v_\infty(z)} \frac{z^{-1}}{s^{-v_\infty(z)}}$$

which is a unit in $\mathcal{O}_\infty$, $\bar{c}$ is the residue class of c in $\mathcal{O}_\infty/s\mathcal{O}_\infty \cong \mathbb{F}_p$, and $\left(\frac{\bar{c}}{p} \right)$ is the Legendre symbol of $\bar{c}$. Note that our formula for c is the reciprocal of the formula in [17] because the reciprocity map in [17] maps uniformizers in K to arithmetic Frobenius elements in $\text{Gal}(\overline{K}_\infty/K_\infty)^{\text{ab}}$, whereas the reciprocity map in [11] maps uniformizers in K to geometric Frobenius elements. One can verify $a(\chi') = 1$. For any $z \in s\mathcal{O}_\infty^*$, write

$$z = s(r_0 + r_1 s + \cdots)$$

with $r_i \in \mathbb{F}_p$ and $r_0 \neq 0$. We then have

$$\begin{aligned} \bar{c} &= -r_0^{-1}, \\ \text{Res}_\infty(z dt) &= -r_0. \end{aligned}$$

So we have

$$\begin{aligned}
\epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) &= \int_{s\mathcal{O}_\infty^*} \chi'^{-1}(z) \psi(\text{Res}_\infty(z dt)) dz \\
&= \int_{s\mathcal{O}_\infty^*} \chi\left(\left(\frac{-r_0^{-1}}{p}\right)\right) \psi(-r_0) dz \\
&= \int_{s\mathcal{O}_\infty^*} \left(\frac{-r_0}{p}\right) \psi(-r_0) dz \\
&= \sum_{r_0 \in \mathbb{F}_p^*} \int_{r_0 s(1+s\mathcal{O}_\infty)} \left(\frac{-r_0}{p}\right) \psi(-r_0) dz \\
&= \sum_{r_0 \in \mathbb{F}_p^*} \left(\frac{-r_0}{p}\right) \psi(-r_0) \int_{r_0 s(1+s\mathcal{O}_\infty)} dz \\
&= \frac{1}{p^2} \sum_{r_0 \in \mathbb{F}_p^*} \left(\frac{-r_0}{p}\right) \psi(-r_0) \\
&= -\frac{g(\chi, \psi)}{p^2}.
\end{aligned}$$

(iv) We can use the same method as in (iii), or use the formula [11] 3.1.5.5 to get

$$\begin{aligned}
\epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}) &= \epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, 2t dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \chi' \left(\frac{2}{s}\right) p^{v_\infty(\frac{2}{s})} \epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \left(\frac{-2}{p}\right) \cdot \frac{1}{p} \cdot \epsilon(\mathbb{P}_{(\infty)}^1, j_* \mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= -\frac{g(\chi, \psi)}{p^3} \left(\frac{-2}{p}\right)
\end{aligned}$$

(v) Taking a to be $at = \frac{a}{s}$ and $b = z$ in the explicit reciprocity law in [17] XIV §5 Proposition 15, we see the character

$$K_\infty^* \rightarrow \overline{\mathbb{Q}_l^*}$$

corresponding to $\mathcal{L}_\chi(at)$ is

$$z \mapsto \psi^{-1} \left(-\text{Res}_\infty \left(\frac{a}{s} \cdot \frac{dz}{z} \right) \right).$$

(We add the negative sign to the formula in [17] since the reciprocity map in [17] is different from the one used in [11].) So the character

$$\phi : K_\infty^* \rightarrow \overline{\mathbb{Q}_l^*}$$

corresponding to $\mathcal{L}_\chi(at) \otimes \mathcal{L}_\chi$ is given by

$$\phi(z) = \psi^{-1} \left(-\text{Res}_\infty \left(\frac{a}{s} \cdot \frac{dz}{z} \right) \right) \chi^{-1} \left(\left(\frac{\bar{c}}{p} \right) \right),$$

where $c = (-1)^{-v_\infty(z)} \frac{z^{-1}}{s^{-v_\infty(z)}}$. One can verify $a(\phi) = 2$. For any $z \in s\mathcal{O}_\infty^*$, write

$$z = s(r_0 + r_1s + \cdots)$$

with $r_i \in \mathbb{F}_p$ and $r_0 \neq 0$. We then have

$$\begin{aligned} \text{Res}_\infty \left(\frac{a}{s} \cdot \frac{dz}{z} \right) &= \frac{ar_1}{r_0}, \\ \bar{c} &= -r_0^{-1}, \\ \text{Res}_\infty(z dt^2) &= -2r_1. \end{aligned}$$

So we have

$$\begin{aligned} &\epsilon(\mathbb{P}_{(\infty)}^1, j_*\mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}) \\ &= \int_{s\mathcal{O}_\infty^*} \phi^{-1}(z) \psi(\text{Res}_\infty(z dt^2)) dz \\ &= \int_{s\mathcal{O}_\infty^*} \psi\left(-\frac{ar_1}{r_0}\right) \chi\left(\left(\frac{-r_0^{-1}}{p}\right)\right) \psi(-2r_1) dz \\ &= \int_{s\mathcal{O}_\infty^*} \left(\frac{-r_0}{p}\right) \psi\left(-r_1\left(\frac{a}{r_0} + 2\right)\right) dz \\ &= \sum_{r_0, r_1 \in \mathbb{F}_p, r_0 \neq 0} \int_{s(r_0 + r_1s)(1+s^2\mathcal{O}_\infty)} \left(\frac{-r_0}{p}\right) \psi\left(-r_1\left(\frac{a}{r_0} + 2\right)\right) dz \\ &= \sum_{r_0, r_1 \in \mathbb{F}_p, r_0 \neq 0} \left(\frac{-r_0}{p}\right) \psi\left(-r_1\left(\frac{a}{r_0} + 2\right)\right) \int_{s(r_0 + r_1s)(1+s^2\mathcal{O}_\infty)} dz \\ &= \frac{1}{p^3} \sum_{r_0, r_1 \in \mathbb{F}_p, r_0 \neq 0} \left(\frac{-r_0}{p}\right) \psi\left(-r_1\left(\frac{a}{r_0} + 2\right)\right) \\ &= \frac{1}{p^3} \sum_{r_0 \in \mathbb{F}_p^*} \left(\frac{-r_0}{p}\right) \sum_{r_1 \in \mathbb{F}_p} \psi\left(-r_1\left(\frac{a}{r_0} + 2\right)\right) \\ &= \frac{1}{p^3} \cdot \left(\frac{-\frac{a}{2}}{p}\right) \cdot p \\ &= \frac{1}{p^2} \left(\frac{2a}{p}\right). \end{aligned}$$

We omit the proof of (vi), which is similar to the proof of (v).

(vii) We have $[2]_*\overline{\mathbb{Q}}_\ell \cong \overline{\mathbb{Q}}_\ell \oplus j_*\mathcal{L}_\chi$. So

$$\epsilon(\mathbb{P}_{(\infty)}^1, [2]_*\overline{\mathbb{Q}}_\ell|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = \epsilon(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_\ell|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \epsilon(\mathbb{P}_{(\infty)}^1, j_*\mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}).$$

We then use (i) and (iii).

(viii) We can define ϵ -factors for virtual sheaves on $\mathbb{P}_{(\infty)}^1$. By [11] 3.1.5.4 (iv), we have

$$\epsilon(\mathbb{P}_{(\infty)}^1, [2]_*([j_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)] - [\overline{\mathbb{Q}}_\ell])|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) = \epsilon(\mathbb{P}_{(\infty)}^1, ([j_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)] - [\overline{\mathbb{Q}}_\ell])|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1}).$$

Hence

$$\begin{aligned} & \epsilon(\mathbb{P}_{(\infty)}^1, j_*[2]_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\ &= \frac{\epsilon(\mathbb{P}_{(\infty)}^1, j_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)|_{\mathbb{P}_{(\infty)}^1}, dt^2|_{\mathbb{P}_{(\infty)}^1})}{\epsilon(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_\ell, dt^2|_{\mathbb{P}_{(\infty)}^1})} \epsilon(\mathbb{P}_{(\infty)}^1, [2]_*\overline{\mathbb{Q}}_\ell, dt|_{\mathbb{P}_{(\infty)}^1}). \end{aligned}$$

We then apply the formulas (ii), (v), and (vii).

We omit the proof of (ix) and (x), which is similar to the proof of (viii). $\square$

Lemma 2.6. *We have*

$$\begin{aligned} & \epsilon(\mathbb{P}_{(\infty)}^1, j_*(\mathcal{L}_{\chi^r} \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^r})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\ &= \begin{cases} \frac{g(\chi, \psi)^{-4r}}{p^2} & \text{if } r \text{ is even,} \\ -\frac{g(\chi, \psi)^{-2r+1}}{p^2} \left(\frac{-1}{p}\right) & \text{if } r \text{ is odd,} \end{cases} \\ & \epsilon(\mathbb{P}_{(\infty)}^1, j_*([2]_*\mathcal{L}_\psi((4i-4r)t) \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^i})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\ &= \begin{cases} -\frac{g(\chi, \psi)^{-6r+1}}{p^4} \left(\frac{-1}{p}\right)^i & \text{if } p|i-r, \\ -\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p}\right)^i & \text{if } p \nmid i-r, \end{cases} \\ & \epsilon(\mathbb{P}_{(\infty)}^1, j_*([2]_*\mathcal{L}_\psi((4i-4r-2)t) \otimes \mathcal{L}_\chi) \otimes \mathcal{L}_{\theta_0^{2r+1}\theta_1^{i+1}}|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\ &= \begin{cases} \frac{g(\chi, \psi)^{-4r}}{p^4} \left(\frac{-2}{p}\right) & \text{if } p|2i-2r-1, \\ -\frac{g(\chi, \psi)^{-2r}}{p^3} \left(\frac{(-1)^{i+1}(2i-2r-1)}{p}\right) & \text{if } p \nmid 2i-2r-1. \end{cases} \end{aligned}$$

Proof. Let F_∞ be the geometric Frobenius element at ∞ . We have

$$\theta_0(F_\infty) = g(\chi, \psi), \quad \theta_1(F_\infty) = \left(\frac{-1}{p}\right).$$

Using the notation [11] 3.1.5.1, we have

$$\begin{aligned} a(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_\ell, dt|_{\mathbb{P}_{(\infty)}^1}) &= -2, \\ a(\mathbb{P}_{(\infty)}^1, j_*\mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) &= -1, \\ a(\mathbb{P}_{(\infty)}^1, [2]_*\overline{\mathbb{Q}}_\ell, dt|_{\mathbb{P}_{(\infty)}^1}) &= -3, \\ a(\mathbb{P}_{(\infty)}^1, j_*[2]_*\mathcal{L}_\psi(at)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) &= -1, \quad (a \in \mathbb{F}_p^*) \\ a(\mathbb{P}_{(\infty)}^1, j_*[2]_*\mathcal{L}_\chi|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) &= -2, \\ a(\mathbb{P}_{(\infty)}^1, j_*[2]_*(\mathcal{L}_\psi(at) \otimes \mathcal{L}_\chi)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) &= -1, \quad (a \in \mathbb{F}_p^*). \end{aligned}$$

So by [11] 3.1.5.6, we have

$$\begin{aligned}
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*(\mathcal{L}_{\chi^r} \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^r})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \begin{cases} ((\theta_0^{2r}\theta_1^r)(F_{\infty}))^{-2}\epsilon(\mathbb{P}_{(\infty)}^1, \overline{\mathbb{Q}}_{\ell}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } r \text{ is even,} \\ ((\theta_0^{2r}\theta_1^r)(F_{\infty}))^{-1}\epsilon(\mathbb{P}_{(\infty)}^1, j_*\mathcal{L}_{\chi}|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } r \text{ is odd,} \end{cases} \\
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*([2]_*\mathcal{L}_{\psi}((4i-4r)t) \otimes \mathcal{L}_{\theta_0^{2r}\theta_1^i})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \begin{cases} ((\theta_0^{2r}\theta_1^i)(F_{\infty}))^{-3}\epsilon(\mathbb{P}_{(\infty)}^1, [2]_*\overline{\mathbb{Q}}_{\ell}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } p|i-r, \\ ((\theta_0^{2r}\theta_1^i)(F_{\infty}))^{-1}\epsilon(\mathbb{P}_{(\infty)}^1, j_*[2]_*\mathcal{L}_{\psi}((4i-4r)t)|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } p \nmid i-r, \end{cases} \\
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*([2]_*(\mathcal{L}_{\psi}((4i-4r-2)t) \otimes \mathcal{L}_{\chi}) \otimes \mathcal{L}_{\theta_0^{2r+1}\theta_1^{i+1}})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \begin{cases} ((\theta_0^{2r+1}\theta_1^{i+1})(F_{\infty}))^{-2}\epsilon(\mathbb{P}_{(\infty)}^1, j_*[2]_*\mathcal{L}_{\chi}|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } p|2i-2r-1, \\ ((\theta_0^{2r+1}\theta_1^{i+1})(F_{\infty}))^{-1}\epsilon(\mathbb{P}_{(\infty)}^1, j_*[2]_*(\mathcal{L}_{\psi}((4i-4r-2)t) \otimes \mathcal{L}_{\chi})|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) & \text{if } p \nmid 2i-2r-1. \end{cases}
\end{aligned}$$

We then apply the formulas in Lemma 2.5. $\square$

The following is Proposition 0.4 in the introduction.

Proposition 2.7. $\epsilon(\mathbb{P}_{(\infty)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1})$ equals

$$p^{-(k+1)(\frac{k+8}{4} + [\frac{k}{2p}])}$$

if $k = 2r$ for an even r ,

$$p^{-(k+1)(\frac{k+6}{4} + [\frac{k}{2p}])}$$

if $k = 2r$ for an odd r , and

$$(-1)^{\frac{k+1}{2} + [\frac{k}{p}] - [\frac{k}{2p}]} p^{-\frac{k+1}{2}(\frac{k+5}{2} + [\frac{k}{p}] - [\frac{k}{2p}])} \left(\frac{-2}{p}\right)^{[\frac{k}{p}] - [\frac{k}{2p}]} \prod_{j \in \{0, 1, \dots, [\frac{k}{2}]\}, p \nmid 2j+1} \left(\frac{(-1)^j(2j+1)}{p}\right)$$

if $k = 2r + 1$.

Proof. By Lemmas 2.4 and 2.6, $\epsilon(\mathbb{P}_{(\infty)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1})$ equals

$$\frac{g(\chi, \psi)^{-4r}}{p^2} \prod_{i \in \{0, \dots, r-1\}, p|i-r} \left(-\frac{g(\chi, \psi)^{-6r+1}}{p^4} \left(\frac{-1}{p}\right)^i\right) \prod_{i \in \{0, \dots, r-1\}, p \nmid i-r} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p}\right)^i\right)$$

if $k = 2r$ for an even r ,

$$-\frac{g(\chi, \psi)^{-2r+1}}{p^2} \left(\frac{-1}{p}\right) \prod_{i \in \{0, \dots, r-1\}, p|i-r} \left(-\frac{g(\chi, \psi)^{-6r+1}}{p^4} \left(\frac{-1}{p}\right)^i\right) \prod_{i \in \{0, \dots, r-1\}, p \nmid i-r} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p}\right)^i\right)$$

if $k = 2r$ for an odd r , and

$$\prod_{i \in \{0, \dots, r\}, p|2i-2r-1} \left(\frac{g(\chi, \psi)^{-4r}}{p^4} \left(\frac{-2}{p}\right)\right) \prod_{i \in \{0, \dots, r\}, p \nmid 2i-2r-1} \left(-\frac{g(\chi, \psi)^{-2r}}{p^3} \left(\frac{(-1)^{i+1}(2i-2r-1)}{p}\right)\right)$$

if $k = 2r + 1$. Let's simplify the above expressions. Recall that $g(\chi, \psi)^2 = p \left(\frac{-1}{p} \right)$. If $k = 2r$ with r even, we have

$$\begin{aligned}
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \frac{g(\chi, \psi)^{-4r}}{p^2} \prod_{i \in \{0, \dots, r-1\}, p|i-r} \left(-\frac{g(\chi, \psi)^{-6r+1}}{p^4} \left(\frac{-1}{p} \right)^i \right) \prod_{i \in \{0, \dots, r-1\}, p \nmid i-r} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p} \right)^i \right) \\
&= \frac{g(\chi, \psi)^{-4r}}{p^2} \prod_{i \in \{0, \dots, r-1\}, p|i-r} \frac{g(\chi, \psi)^{-4r}}{p} \prod_{i \in \{0, \dots, r-1\}} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p} \right)^i \right) \\
&= \frac{g(\chi, \psi)^{-4r}}{p^2} \left(\frac{g(\chi, \psi)^{-4r}}{p} \right)^{\left[\frac{r}{p}\right]} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \right)^r \left(\frac{-1}{p} \right)^{\frac{r(r-1)}{2}} \\
&= \frac{p^{-2r}}{p^2} \left(\frac{p^{-2r}}{p} \right)^{\left[\frac{r}{p}\right]} \frac{\left(p \left(\frac{-1}{p} \right) \right)^{\frac{r(-2r+1)}{2}}}{p^{3r}} \left(\frac{-1}{p} \right)^{\frac{r(r-1)}{2}} \\
&= p^{-r^2 - \frac{9}{2}r - 2 - (2r+1)\left[\frac{r}{p}\right]} \left(\frac{-1}{p} \right)^{-\frac{r^2}{2}} \\
&= p^{-(k+1)\left(\frac{k+8}{4} + \left[\frac{k}{2p}\right]\right)}.
\end{aligned}$$

If $k = 2r$ with r odd, we have

$$\begin{aligned}
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= -\frac{g(\chi, \psi)^{-2r+1}}{p^2} \left(\frac{-1}{p} \right) \prod_{i \in \{0, \dots, r-1\}, p|i-r} \left(-\frac{g(\chi, \psi)^{-6r+1}}{p^4} \left(\frac{-1}{p} \right)^i \right) \\
&\quad \times \prod_{i \in \{0, \dots, r-1\}, p \nmid i-r} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p} \right)^i \right) \\
&= -\frac{g(\chi, \psi)^{-2r+1}}{p^2} \left(\frac{-1}{p} \right) \prod_{i \in \{0, \dots, r-1\}, p|i-r} \left(\frac{g(\chi, \psi)^{-4r}}{p} \right) \prod_{i \in \{0, \dots, r-1\}} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \left(\frac{-1}{p} \right)^i \right) \\
&= -\frac{g(\chi, \psi)^{-2r+1}}{p^2} \left(\frac{-1}{p} \right) \left(\frac{g(\chi, \psi)^{-4r}}{p} \right)^{\left[\frac{r}{p}\right]} \left(-\frac{g(\chi, \psi)^{-2r+1}}{p^3} \right)^r \left(\frac{-1}{p} \right)^{\frac{r(r-1)}{2}} \\
&= \frac{g(\chi, \psi)^{(-2r+1)(r+1)}}{p^{3r+2}} \left(\frac{-1}{p} \right)^{1 + \frac{r(r-1)}{2}} \left(\frac{g(\chi, \psi)^{-4r}}{p} \right)^{\left[\frac{r}{p}\right]} \\
&= \frac{\left(p \left(\frac{-1}{p} \right) \right)^{\frac{(-2r+1)(r+1)}{2}}}{p^{3r+2}} \left(\frac{-1}{p} \right)^{1 + \frac{r(r-1)}{2}} \left(\frac{p^{-2r}}{p} \right)^{\left[\frac{r}{p}\right]} \\
&= p^{-r^2 - \frac{7}{2}r - \frac{3}{2} - (2r+1)\left[\frac{r}{p}\right]} \left(\frac{-1}{p} \right)^{-\frac{(r-1)(r+3)}{2}} \\
&= p^{-(k+1)\left(\frac{k+6}{4} + \left[\frac{k}{2p}\right]\right)}.
\end{aligned}$$

If $k = 2r + 1$ is odd, we have

$$\begin{aligned}
& \epsilon(\mathbb{P}_{(\infty)}^1, j_*(\text{Sym}^k(\text{Kl}_2))|_{\mathbb{P}_{(\infty)}^1}, dt|_{\mathbb{P}_{(\infty)}^1}) \\
&= \prod_{i \in \{0, \dots, r\}, p \nmid 2i-2r-1} \left(\frac{g(\chi, \psi)^{-4r}}{p^4} \left(\frac{-2}{p} \right) \right) \prod_{i \in \{0, \dots, r\}, p \nmid 2i-2r-1} \left(-\frac{g(\chi, \psi)^{-2r}}{p^3} \left(\frac{(-1)^{i+1}(2i-2r-1)}{p} \right) \right) \\
&= \left(-\frac{g(\chi, \psi)^{-2r}}{p^3} \right)^{r+1} \left(-\frac{g(\chi, \psi)^{-2r}}{p} \left(\frac{-2}{p} \right) \right)^{\left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} \prod_{i \in \{0, \dots, r\}, p \nmid 2i-2r-1} \left(\frac{(-1)^{i+1}(2i-2r-1)}{p} \right) \\
&= \left(-p^{-3-r} \left(\frac{-1}{p} \right)^{-r} \right)^{r+1} \left(-p^{-1-r} \left(\frac{-1}{p} \right)^{-r} \left(\frac{-2}{p} \right) \right)^{\left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} \prod_{j \in \{0, \dots, r\}, p \nmid 2j+1} \left(\frac{(-1)^{r-j}(2j+1)}{p} \right) \\
&= (-1)^{r+1 + \left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} p^{-(r+1)(r+3 + \left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right])} \left(\frac{-2}{p} \right)^{\left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} \left(\frac{-1}{p} \right)^{-r(r+1) - r(\left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right])} \\
&\quad \times \prod_{j \in \{0, \dots, r\}, p \nmid 2j+1} \left(\frac{(-1)^{r-j}(2j+1)}{p} \right) \\
&= (-1)^{\frac{k+1}{2} + \left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} p^{-\frac{k+1}{2}(r+3 + \left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right])} \left(\frac{-2}{p} \right)^{\left[\frac{k}{p}\right] - \left[\frac{k}{2p}\right]} \prod_{j \in \{0, \dots, \left[\frac{k}{2}\right]\}, p \nmid 2j+1} \left(\frac{(-1)^j(2j+1)}{p} \right).
\end{aligned}$$

□

References

- [1] M. Artin, A. Grothendieck, and J.-L. Verdier, Théorie des topos et cohomologie étale des schémas (SGA 4), Lecture Notes in Math. 269, 270, 305, Springer-Verlag (1972-1973).
- [2] H.T. Choi and R.J. Evans, Congruences for sums of powers of Kloosterman sums, Inter. J. Number Theory, 3(2007), 105-117.
- [3] P. Deligne, Applications de la Formule des Traces aux Sommes Trigonométriques, in Cohomologie Étale (SGA 4 $\frac{1}{2}$), 168-232, Lecture Notes in Math. 569, Springer-Verlag 1977.
- [4] R. J. Evans, Seventh power moments of Kloosterman sums, Israel J. Math., to appear.
- [5] L. Fu, Calculation of ℓ -adic local Fourier transformations, arXiv: 0702436 [math.AG] (2007).
- [6] L. Fu and D. Wan, Trivial factors for L -functions of symmetric products of Kloosterman sheaves, Finite Fields and Appl. 14 (2008), 549-570.
- [7] L. Fu and D. Wan, L -functions for symmetric products of Kloosterman sums, J. Reine Angew. Math. 589 (2005), 79-103.

- [8] L. Fu and D. Wan, L -functions of symmetric products of the Kloosterman sheaf over $\mathbb{Z}$, Math. Ann. 342 (2008), 387-404.
- [9] K. Hulek, J. Spandaw, B. van Geemen and D. van Straten, The modularity of the Barth-Nieto quintic and its relative, Adv. Geom., 1(2001), no.3, 263-289.
- [10] N. Katz, Gauss sums, Kloosterman sums, and monodromy groups, Princeton University Press 1988.
- [11] G. Laumon, Transformation de Fourier, constantes d'équations fonctionnelles, et conjecture de Weil, Publ. Math. IHES 65 (1987), 131-210.
- [12] M. Moisio, The moments of Kloosterman sums and the weight distribution of a Zetterberg-type binary cyclic code, IEEE, Trans. Inform. Theory, 53(2007), 843-847.
- [13] M. Moisio, On the moments of Kloosterman sums and fibre products of Kloosterman curves, Finite Fields Appl., 14(2008), 515-531.
- [14] C. Peters, J. Top and M. van der Vlugt, The Hasse zeta function of a K3 surface related to the number of words of weight 5 in the Melas codes, J. Reine Angew. Math. 432(1992), 151-176.
- [15] P. Robba, Symmetric powers of p -adic Bessel functions, J. Reine Angew. Math., 366(1986), 194-220.
- [16] J.-P. Serre, Linear representations of finite groups, Springer-Verlag (1977).
- [17] J.-P. Serre, Local fields, Springer-Verlag, 1979.
- [18] D. Wan, Dwork's conjecture on unit root zeta functions, Ann. Math., 150(1999), 867-927.